



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

mal whorl on the stem, are three more whorls of three leaves each set closely together, making a pretty rosette, and above those is the flower erect. Sepals rather larger than common; petals 11^l long and 4^l wide, with a white stripe running down the center, and a green one on each edge; stigmas four; one of the petals two-parted.

In the other plant a rosette of two whorls; a third abnormal whorl in this specimen also, but lifted half an inch above the others to the base of the flower. Petals green and white; one stamen abortive; stigmas two. The rosettes pressed and mounted measure in one plant 5ⁱⁿ 10^l across, in the other 4ⁱⁿ 11^l.

The upright flowers suggested *T. grandiflorum*, but the stigmas, distinct and recurved, and the short stamens are perhaps enough to prove the species to be *cernuum*; moreover, I learn that *T. grandiflorum* has never been noticed in that vicinity, and that *T. cernuum* is common where these plants were found. They were collected by a very youthful observer, Miss Anna Dimmock, who writes to me that there were several other similar specimens where she gathered those which she sent here. —MARIA L. OWEN, *Springfield, Mass.*

An abnormal *Hepatica*.—A peculiar form of *Hepatica triloba*, collected near Boston, was brought to me recently. The peduncle, which is very flat, though not much larger than usual, is doubtless a case of fasciation. The involucre consists of seven green bracts of unequal size, one with a white streak on both sides. A normal flower of nine semi-white sepals is in the normal position. Another smaller flower, of six white sepals, is crowded between the larger flower and the involucre on a short pedicel. This flower has an involucre of three bracts, one green, and two of a variegated green and white color. Both flowers have the usual number of stamens and pistils. —WALTER DEANE, *Cambridge, Mass.*